## QUANTUM FIELD THEORY, PROBLEM SHEET 3

## Problem 1: The free scalar field and causality

Recall from quantum mechanics that, if two observables are represented by operators  $\mathcal{O}_1$  and  $\mathcal{O}_2$  with  $[\mathcal{O}_1, \mathcal{O}_2] \neq 0$ , then a measurement of  $\mathcal{O}_1$  will influence a subsequent measurement of  $\mathcal{O}_2$ . However, in a Lorentz invariant quantum field theory, two events with spacelike separation should not affect each other in order to preserve causality.

Convince yourself that for any spacelike four-vector z, there exists a proper orthochronous Lorentz transformation sending  $z^0 \to 0$ . Conclude that

$$\Delta(x, y) = 0$$
 whenever  $(x - y)^2 < 0$ .

Here  $\Delta(x,y) = [\phi(x),\phi(y)]$  and  $\phi$  is a free real scalar field. What is the corresponding statement for a complex scalar field?

## Problem 2: The residue theorem

To obtain the electrostatic potential of a point particle in Exercise 3.4 on Problem Sheet 1, you were given the identity

$$\int_0^\infty dk \, \frac{k \, \sin(kx)}{k^2 + m^2} = \frac{\pi}{2} e^{-mx} \qquad (x > 0) \, .$$

Prove this formula with the help of the residue theorem.

## Problem 3: Propagators

You have seen in the lecture that the expression

$$D_{\mathcal{C}}(x-y) = \int_{\mathcal{C}} \frac{d^4k}{(2\pi)^4} \frac{i}{k^2 - m^2} e^{-ik \cdot (x-y)}$$

depends on the curve  $\mathcal C$  in the plane of complex  $k^0$  along which the poles at  $\pm \omega$  are avoided. For example, choosing to circumvent the pole at  $k^0=-\omega$  in the lower half-plane and the pole at  $k^0=+\omega$  in the upper half-plane yields the Feynman propagator

$$D_F(x-y) = \int \frac{d^4k}{(2\pi)^4} \frac{i}{k^2 - m^2 + i\epsilon} e^{-ik\cdot(x-y)} = \langle 0|T \phi(x)\phi(y)|0\rangle$$
$$= \Theta(x^0 - y^0)\langle 0|\phi(x)\phi(y)|0\rangle + \Theta(y^0 - x^0)\langle 0|\phi(y)\phi(x)|0\rangle.$$

1. Show that any  $iD_{\mathcal{C}}(x-y)$  is a Green function for the Klein-Gordon operator  $\Box + m^2$ :

$$(\Box_x + m^2)D_{\mathcal{C}}(x - y) = -i\delta^4(x - y).$$

- 2. Express  $D_R(x-y)$  and  $D_A(x-y)$  in terms of  $\Theta$  functions and of vacuum expectation values of products of  $\phi(x)$  and  $\phi(y)$ . Here  $D_R(x-y)$  is defined by a contour which circumvents both poles in the upper half-plane, and  $D_A(x-y)$  by a contour which circumvents both poles in the lower half-plane.
- 3. Starting from the expression of the lecture

$$D_F(x-y) = \int \widetilde{dk} \left( e^{-ik(x-y)} \Theta(x^0 - y^0) + e^{ik(x-y)} \Theta(y^0 - x^0) \right) \Big|_{k^0 = \omega_{\vec{k}}}$$

and evaluating the integral, write the Feynman propagator for  $(x - y)^2 \neq 0$  explicitly in terms of the modified Bessel function of the second kind  $K_1(z)$ . You can use the identity (see Gradshteyn & Ryzhik, "Table of integrals, series and products", eq. 3.914/9)

$$\int_0^\infty \frac{xe^{-\beta\sqrt{\gamma^2+x^2}}}{\sqrt{\gamma^2+x^2}}\sin bx\,\mathrm{d}x = \frac{\gamma b}{\sqrt{\beta^2+b^2}}K_1\left(\gamma\sqrt{\beta^2+b^2}\right)\,.$$

For illustration, this is how  $K_1$  behaves for real and imaginary arguments:



